



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.



Disponible en ligne sur

ScienceDirect
www.sciencedirect.com

Elsevier Masson France

EM|consulte
www.em-consulte.com



CORONAVIRUS: OPINION OF THE FRENCH NATIONAL ACADEMY OF MEDICINE

Vitamin D and COVID-19[☆]

Available online 30 May 2020

Vitamin D is a prohormone synthesized in the dermis under the effect of ultraviolet light, i.e. sunrays, and then transported to the liver and kidney, where it is transformed into an active hormone. It is responsible for intestinal absorption of calcium and bone health.

But vitamin D also has unconventional effects. In particular, it modulates the functioning of the immune system by stimulating macrophages and dendritic cells [1–3]. It plays a role in regulating and suppressing the cytokine inflammatory response that causes the acute respiratory distress syndrome that characterizes the severe and often lethal forms of COVID-19.

A significant correlation between low serum vitamin D levels and mortality from the COVID-19 has been shown [4]. This phenomenon generally follows a north-south gradient, although there are exceptions such as the Nordic countries, where vitamin D nutrient supplementation, particularly from milk products, is systematic. However, Southern European countries show a surprisingly high prevalence of vitamin D deficiency, despite higher levels of sunlight [5]. This would explain why infants, who receive vitamin D regularly, have asymptomatic forms of COVID-19 and fewer complications.

Vitamin D cannot be considered as a preventive or a curative treatment for CoV-2 SARS infection. However, by mitigating the inflammatory storm and its consequences, it could be considered as an adjunct to any form of therapy.

The National Academy of Medicine:

- points out that the administration of vitamin D per os is a simple and inexpensive measure, that is reimbursed by the French National Health Insurance;
- confirms its recommendation to ensure vitamin D supplementation in the French population in a 2012 report [2];
- recommends the rapid serum vitamin D (i.e. 25 OHD) testing in people over 60 years of age with COVID-19, and a loading dose of 50,000 to 100,000 IU in case of deficiency, which could help limit respiratory complications;
- recommends vitamin D supplementation of 800 to 1000 IU/day in people under 60, as soon as the diagnosis of COVID-19 is confirmed.

Disclosure of interest

The authors declare that they have no competing interest.

References

- [1] Liu PT, Stenger S, et al. Toll like receptor triggering of a vitamin D mediated human antimicrobial response. *Science* 2006;311:1770.
- [2] Salle B, au nom d'un groupe de travail de l'Académie nationale de médecine. Statut vitaminique, rôle extra-osseux et besoins quotidiens en vitamine D. *Bull Acad Natl Med* 2012;196:1011–5.
- [3] Laird E, Rhodes JM, Kenny RA. Vitamin D and inflammation: potential implications for severity of COVID-19. *Irish med J* 2020;113:81.
- [4] McCartney DM, Byrne DG. Optimisation of vitamin D status impact mortality from SARS-CoV-2 infection. *Irish Med J* 2020;113:58.
- [5] Lips P, Cashman KD, et al. Current vitamin D status in European and Middle East countries and strategies to prevent vitamin D deficiency: a position statement of the European Calcified Tissue Society. *Eur J Endocrinol* 2019;180:23–54.

DOI of original article: <https://doi.org/10.1016/j.banm.2020.05.097>.

[☆] Press release from the French National Academy of Medicine, 22 May 2020.

<https://doi.org/10.1016/j.banm.2020.05.098>
0001-4079/